

Dissertation Summary

Stellar populations in local star-forming galaxies

Pablo G. Pérez-González

Current address: University of Arizona. Steward Observatory. 933 N Cherry Av. Tucson,
AZ85721, USA

Electronic mail: pgperez@as.arizona.edu

Thesis work conducted at: Departamento de Astrofísica, Facultad de Ciencias Físicas,
Universidad Complutense de Madrid (Spain)

Ph.D. Thesis directed by: J. Zamorano & J. Gallego ; Ph.D. Degree awarded: June 2003

Received _____

Subject headings: astronomical databases: surveys, galaxies: fundamental
parameters, galaxies: luminosity function, mass function, galaxies: photometry,
galaxies: starburst, galaxies: stellar content

One of the main issues in today's Astrophysics is how present day galaxies formed and how they have evolved over time. A considerable observational effort is being made to study galaxies from the earliest possible times to the present. One approach to understand how present-day galaxies came into being is to study in detail the properties of local active star-forming galaxies, and in particular their star-formation histories. In this respect, it is important to quantify the relative importance of the current episode of star formation in comparison to the underlying older stellar populations.

The main goal of this thesis work ¹ is studying the main properties of the stellar populations embedded in a statistically complete sample of local active star-forming galaxies: the Universidad Complutense de Madrid (UCM) Survey of emission-line galaxies ². This sample contains 191 local star-forming galaxies at an average redshift of 0.026. The survey was carried out using an objective-prism technique centered at the wavelength of the H α nebular emission-line (a common tracer of recent star formation).

We have compiled an extensive multi-wavelength (from the optical to the near infrared) photometric (in narrow and broad band filters) and spectroscopic dataset for all the galaxies in the sample. The analysis of this dataset has been centered on the characterization of the morphology of the objects, on the comparison of the global properties of the sample with those of other galaxy samples at several redshifts, and on the study of the stellar content of the UCM galaxies. In the latter research topic, a special attention has been paid to the most recent star-forming bursts, which are the main characteristic of our objects. We have also developed (and tested with the dataset for the UCM sample) a powerful stellar population synthesis tool (which is able to obtain robust estimates of the stellar mass, a very important parameter in the study of the evolution of galaxies), and a technique to calculate luminosity and stellar mass functions. These techniques may be easily used in the characterization of samples of galaxies at any redshift.

The main results obtained in this work point out that the UCM Survey galaxies span a broad range in properties (among others, the total stellar mass and the star formation rate per stellar mass unit) between those of galaxies completely dominated by

¹The original text is written in Spanish and is available at http://t-rex.fis.ucm.es/Publications/Repository/tesis_pag_1.ps.gz and http://t-rex.fis.ucm.es/Publications/Repository/tesis_pag_2.ps.gz

²http://t-rex.fis.ucm.es/UCM_Survey/index.html

current/recent star formation (e.g., extreme dwarf HII galaxies or starburst galaxies) and those of ‘normal’ quiescent spirals. The UCM objects present intermediate-late Hubble type spiral morphologies and a recent instantaneous burst of star formation (occurred about 5 Myr ago) with sub-solar metallicity. These bursts involved $\sim 5\%$ of the total stellar mass of the UCM galaxies. An ‘average’ UCM galaxy has a total stellar mass of 10^{10} solar masses, i.e., about a factor of 7-10 lower than a typical local spiral galaxy. Finally, we estimate that $13 \pm 3\%$ of the total baryon mass density in the form of stars in the local Universe is associated with star-forming galaxies.